



Effects of certain seaweed extracts on the primary bio-film forming bacteria

¹V. Balasubramanian, ²S. Palanichamy and ^{*1}R. Rajaram

¹Department of Marine Science, Bharathidasan University, Tiruchirappalli, Tamil Nadu - 620 024.

*E-mail: rrajaram69@rediffmail.com

²Offshore Platform and Marine Electrochemistry Centre, CECRI unit, Harbour area, Tuticorin, Tamil Nadu - 628 004, India

Abstract

The present investigation is an attempt to bring out the specific antifouling properties of extracts of five species of marine algae collected from the Gulf of Mannar area of Tuticorin coast. The extracts from seaweeds namely, *Ulva lactuca*, *Caulerpa scalpelliformis*, *Padina boergeresii*, *Caulerpa* sp. and *Chaetomorpha linoides* were tested against biofilm forming bacteria namely, *Micrococci* sp., *Aeromonas* sp., *Pseudomonas* sp., *Flavobacterium* sp., *Cytophaga* sp. and *Enterobacter* sp. The disc diffusion method was used to find out the inhibition of biofilm forming bacteria through zone of inhibition. Among five algal genera, four species were found to have antagonistic activity against biofilm forming bacterial isolates. We conclude that the broad antifouling properties of the crude extracts of the four seaweed species inhibit the growth of biofilm forming bacteria.

Keywords: Seaweeds, antifouling, biofilm bacteria

Introduction

Marine biofouling is a phenomenon causing large penalties to engineered structures such as ships and offshore platforms by way of increased use of manpower, fuel, material and dry-docking time (Chambers *et al.*, 2006). Materials immersed in seawater are acted upon by a series of physical, chemical and biological events which result in the formation of a biofilm complex, including a layer of attached organisms like bacteria, fungi, algae, diatoms and barnacles and the colonization of biofoulants depends on polluted nature of the environment (Karande, 1968, 1978; Charaklis, 1981; Karande and Srivastava, 1984; Srivastava *et al.*, 1990; Abarzua and Jakubowski, 1995). Immediately after immersion surfaces are coated with a glycoproteinaceous film which favors the colonisation by bacteria, fungi, diatom, protozoa and other microorganisms. Biofilms are clusters of microbial cells that are attached to a surface. They occur in nearly every moist environment where sufficient nutrient flow is available and surface attachment can be achieved. A biofilm can be formed

by a single bacterial species, although they also consist of many species of bacteria, fungi, algae and protozoa. Antifouling compounds produced by bacteria are the main components of biofilms in a marine environment and possibly show important effects on biofouling (Maki, 2002). Bacteria in a biofilm can also affect the growth of other bacteria in the same biofilm (Burgess *et al.*, 1999). For example, the presence of “resident” bacterial strains on particles either increases or decreases the colonisation rate of “newcomer” strains (Grossart *et al.*, 2003). Macrophytes and invertebrates such as barnacles and blue mussels subsequently attach to this film (Davis *et al.*, 1989). The control of biofouling is of particular concern in modern marine engineering and shipping operations and is one of the most important problems currently facing marine technology (Hattori and Shizui, 1996). Uncontrolled settlement and adhesion of marine invertebrates and algae to the hulls of ships increase frictional drag, with a corresponding decrease in speed, maneuverability and fuel efficiency.

Marine algae play an important role in the fouling

of a wide range of immersed artificial substrata, particularly in shallow water where there is sufficient light to permit the active growth of algae. Although there are many diverse phyla of marine algae, only relatively small numbers are considered to be important as fouling organisms. The three most economically important phyla are the Chlorophyta, Heteroconta and Rhodophyta; Cyanophyta are also frequently reported particularly as primary colonizers (Fletcher, 1980). Microalgae and macro algae such as *Enteromorpha intestinalis* and *Ulothrix zonata* settle after development of a primary film comprising bacteria, with a clear quantitative domain of benthic diatoms (Abarzua and Jakubowski, 1995). Uncontrolled adhesion of fouling by macroalgae such as *Ulva* sp. and *Enteromorpha* sp. cause serious problem by settling on ships, hulls, power plant cooling system (Fletcher, 1980; Miki *et al.*, 1996). Macroalgae are a rich source of natural bioactive products although little has been done to define an ecological role for these compounds. It is therefore possible that they possess chemical defenses to prevent the colonization of the surfaces.

Many substances obtained from marine algae such as alginate, carrageen and agar as phycocolloids have been used for decades (Taskin *et al.*, 2001). Since algae have been used in traditional medicine for a long time (Fitton, 2006) and also some algal substances have bacteriostatic and bactericidal activity, they have been extensively studied by several researchers (Ghosh *et al.*, 2004; Freile - Pelegrín and Morales, 2004; Salvador *et al.*, 2007). Seaweeds are the major principle producers of oceanic plant community, distributed widely and indigenously and are recognised for their elaborate chemical defense characteristics against many biotic and abiotic factors (Aseer *et al.*, 2010). A variety of seaweed biogenic compounds exhibiting antifouling activity, which belongs to the group of fatty acids, lipopeptides, amides, alkaloids, terpenoids, lactones, pyrroles and sterols have been discovered (Bhadury and Wright, 2004). As an alternative, the seaweeds can be utilised for the development of environmentally compatible antifouling agents. The aim of this work is to investigate the antifouling activity of seaweeds collected from Tuticorin coast against the biofilm forming bacteria. Most of the

earlier studies are on the antibacterial activity of seaweed extracts against pathogenic bacteria. There is no study on the antifouling activity of seaweed extract against biofilm forming bacteria. The use of marine natural products that are capable of inhibiting one or several steps in fouling may provide a solution to the marine biofouling. Considering the significance of the issue, five macroalgae were screened for their antifouling activity against biofilm bacteria.

Material and Methods

Collection of seaweeds: Live and healthy samples of the seaweeds *Ulva lactuca*, *Caulerpa scalpelliformis*, *Padina boergesenii*, *Caulerpa* sp. and *Chaetomorpha linoides* were collected by handpicking during low tide from Hare Island in the Gulf of Mannar of Tuticorin coast (08°46'25.15" N lat., 78°11'16.05" E long.).

Preparation of seaweed samples: Seaweeds were thoroughly washed with seawater to remove all epiphytes, shells, barnacles etc. and again washed with freshwater to remove the surface salts, sand particles if any and allowed to air dry at room temperature for 3 to 4 days. Air drying will retain certain amount of moisture; it can be dried in hot air oven at 45-50°C and then pulverised.

Extraction of bioactive principles from seaweeds: For the extraction of crude bioactivities, 20 g of seaweed powder was taken in a 250 ml conical flask. About the same volume of solvents (v/v) like di-ethylether (DEE), chloroform and methanol were added to get the natural concentrations of seaweeds and they were extracted by cold steep method at -10°C (Wright, 1998).

Enumeration of bacteria in biofilm: Biofilm generated on the PVC coupons exposed to natural seawater was scrapped using sterile brush and immediately transferred to sterile saline water and serially diluted using sterile seawater. Samples were inoculated by spread plating method on Zobell Marine Agar to enumerate total aerobic heterotrophic bacteria (THB). The pure cultures were maintained in agar slant at 4°C for bacterial characterisation. The isolate was characterised using various morphological, biochemical and molecular

characters. Gram staining and motility tests were performed for preliminary identification of the isolate (Allegrucci and Sauer, 2007). Morphological parameters include colony color, size, shape and margin studied by microscopic observation. Biochemical tests such as MR-VP, indole, oxidase, catalase, sugar fermentation and nitrate reduction test were carried out (Dalton *et al.*, 1994).

Antibacterial assay: Disc diffusion method (Bauer *et al.*, 1966) was employed for the study of antifouling activity of seaweed extract. Sterile nutrient agar plates were prepared for the purpose. Cotton swabs were autoclaved and used to inoculate the culture on the dried surface of agar plates to ensure the even distribution on the medium. The isolated pure cultures of *Micrococci* sp., *Aeromonas* sp., *Pseudomonas* sp., *Flavobacterium* sp., *Cytophaga* sp. and *Enterobacter* sp. from the biofilm were swabbed on to separate agar plates. After 24–36 hr of incubation at 37°C, the maximum inhibition zone area was 8 mm and minimum was 1 mm.

Results

The results in terms of inhibition zone are given in Fig. 1 to 5. Among the five extracts of seaweeds tested, methanolic extract of *Ulva lactuca* showed considerable antibacterial activity against the *Micrococci* sp. and the maximum zones of inhibition is 8 mm and minimum zone of inhibition was observed in *Cytophaga* sp. (3 mm). The chloroform extract of *Ulva lactuca* showed a maximum zone of inhibition (6 mm) in *Flavobacterium* sp. and *Aeromonas* sp. exhibited a minimal zone of clearance (4 mm). The DEE extract was found to be quite ineffective against most of the biofilm bacteria. However, *Pseudomonas* sp. was found to be sensitive (6.0 mm) while the inhibitory zone was in trace (<1 mm) amount for *Cytophaga* sp (Fig. 1).

The methanolic extracts of *Caulerpa scalpelliformis* showed maximum antibacterial activity against the *Micrococci* sp. (7 mm) and minimum zone of clearance was observed in *Flavobacteria* sp. (2 mm). The chloroform extracts of *C. scalpelliformis* showed a maximal zone of inhibition (6 mm) against *Cytophaga* sp. while *Aeromonas* sp. was found to be rather tolerant, which




Fig. 1. Antibacterial activity of the marine algae *Ulva lactuca*

exhibited only a minimum zone of clearance (1 mm). As observed for *Ulva lactuca*, the DEE extracts of *C. scalpelliformis* were found to be quite ineffective against most of the biofilm bacteria. However, *Cytophaga* sp. was found to be rather sensitive (3 mm) while the inhibitory zone was less amount (1 mm) for all other bacterial strains (Fig. 2).




Fig. 2. Antibacterial activity of the marine algae *Caulerpa scalpelliformis*

The methanolic extract of *Padina boergesenii* showed maximum antibacterial activity against the *Micrococci* sp. (8 mm) and minimum zone of clearance was in *Flavobacterium* sp. (<1 mm). The chloroform extracts of *P. boergesenii* showed a maximal zone of inhibition (7 mm) in *Pseudomonas* sp. while *Micrococci* sp. was found to be tolerant (1 mm). As noted for other seaweeds, the DEE extracts of *P. boergesenii* were found to be quite ineffective against most of the biofilm bacteria. However, *Pseudomonas* sp. was found to be sensitive (5 mm), while inhibitory zone was in trace amount (<1 mm) in all others (Fig. 3).




Fig. 3. Antibacterial activity of the marine algae *Padina boergesenii*

The methanolic extract of *Caulerpa* sp. showed maximum antibacterial activity against *Micrococci* sp., *Pseudomonas* sp. and *Enterobacter* sp. (8 mm), while the zone of clearance was found to be in traces for *Flavobacterium* sp. (1mm). The chloroform extracts of *Caulerpa* sp. showed a maximal zone of inhibition of 7mm for *Pseudomonas* sp. and *Cytophaga* sp., while *Aeromonas* sp. and *Flavobacterium* sp. exhibited a minimal zone of inhibition of 4 mm. The DEE extracts of *Caulerpa* sp. were found to be quite ineffective against most of the biofilm bacteria. However, *Pseudomonas* sp. was found to be sensitive (7 mm), while *Flavobacterium* sp. and *Aeromonas* sp. showed only a minimal zone of 3mm inhibition (Fig. 4).

The methanolic extract of *Chaetomorpha linoides* showed highest antibacterial activity against the *Flavobacterium* sp. (10 mm), while the minimal zone




Fig. 4. Antibacterial activity of the marine algae *Caulerpa* sp.

of clearance was observed in *Enterobacter* sp. (4 mm). Similarly, the chloroform extracts of *C. linoides* showed an appreciable level of inhibition zone against *Flavobacterium* sp. (9 mm) while *Micrococci* sp. was found to be rather tolerant (1 mm). The DEE extracts of *C. linoides* showed no activity against most of the biofilm bacteria. However, *Flavobacterium* sp. was found to be sensitive (5 mm), while *Micrococcus* sp., *Cytophaga* sp., *Aeromonas* sp. and *Enterobacter* sp. were found to have traces (<1 mm) of inhibitory zone (Fig. 5).




Fig. 5. Antibacterial activity of the marine algae *Chaetomorpha linoides*

Discussion

The primary biofilm provides a supporting substrate for subsequent attachment of other fouling organisms. The deterrence of the formation of this layer is fundamental to effective control of further large scale biofouling. The marine macroalgae are well documented as possessing generalised antibacterial and antifungal activity (Caccamese *et al.*, 1980, 1985; Naqvi *et al.*, 1980; Padma Sridher *et al.*, 1984; Pesando and Caram 1984; Bernard and Pesando 1989; Vlachos *et al.*, 1996). Phlorotannins from brown algae have been implicated in antifouling (Sieburth and Conover, 1965; McLachlan and Craigie, 1966; Langlosis, 1975; Fletcher, 1989). Philipps and Towers (1982) suggest that the red algae *Rhodomela larix* may act as an antifoulant. De Nys *et al.* (1995) showed that the red alga *Delisea pulchra* produced a suite of unique secondary metabolites, furanones, that inhibit surface colonization traits in epiphytic marine bacteria without toxicity.

Among the tested biofilm bacteria, *Micrococci* sp. and *Pseudomonas* sp. were found to be sensitive when compared to all other heterotrophs. *Flavobacterium* sp. and *Cytophaga* sp. showed moderate sensitivity, while *Aeromonas* sp. and *Enterobacter* sp. were found to be resistant in some cases. However, from the present study, it is evident that all the five seaweeds possess antifouling compounds. *Ulva* sp., *C. sp.* and *Caulerpa* sp. were found to have remarkable activity, while *Padina* sp. showed mild activity. The degree of antibacterial activities of the seaweed extracts was found to be solvent specific. The algae extracted in methanol and chloroform was found to show considerable antifouling activity, while DEE exhibited poor activity against biofilm bacteria. Sastry and Rao (1994) showed antibacterial activity against both gram positive and gram negative pathogenic bacteria after successive extraction with benzene, chloroform and methanol. Similarly, Mahasneh *et al.* (1995) have shown antibacterial activity in organic extracts of six species of marine algae against multi-antibiotic resistant bacteria.

In the present study, almost all the algal extracts showed antifouling activity against most of the biofilm bacteria tested indicating that these algae are potential sources of antifouling compounds. This may be further investigated with various fractions of the extracts. Caccamese *et al.* (1985) reported that brown and red algal extract showed higher antibacterial activity against human pathogenic bacteria like *Bacillus* sp. and *E. coli*. Similarly, Padmini (1991) reported that red and brown algal extracts showed greater antibacterial activity than green algae. On the contrary, in the present study, green algae were found to be more effective against biofilm forming bacteria compared to brown algae (*Padina* sp.). The variation in the effects of algal extracts suggests that they are not simply functioning as broad spectrum toxins against marine biofilm bacteria; rather they appear to have specific activities against one or several biofilm bacteria.

Acknowledgements

The authors thank the Director, CECRI, Karaikudi for providing facilities and permission for communicating the results. The authors also thank

Dr. S. Maruthamuthu, Scientist, CECRI-Karaikudi for his valuable suggestion and helpful discussions.

References

- Abarzua, S. and S. Jakubowski. 1995. Biotechnological investigation for the prevention of biofouling. I. Biological and biochemical principles for the prevention of biofouling. *Mar. Ecol. Prog. Ser.*, 123: 301–312.
- Allegrucci, M. and K. Sauer. 2007. Characterization of colony morphology variants isolated from *Streptococcus pneumoniae* biofilms. *J. Bacteriol.*, 189(5): 2030–2038.
- Aseer, M., S. Sujith, B. Sabarathnam, G. Seghal Kiran, Joseph Selvin, Chippu Shakir and Aaron Premnath Lipton. 2010. Antifouling potentials of seaweeds collected from the southwest coast of India. *World J. Agricultural Sciences*, 6(3): 243–248.
- Bauer, A. W., W. M. Kirby, J. C. Sherris and M. Turk. 1966. Antibiotic susceptibility by a standardized single disk method. *Am. J. Clin. Pathol.*, 45: 493–496.
- Bernard, P. and D. Pesando. 1989. Antibacterial and antifungal activity of extracts from the rhizomes of the Mediterranean seagrass *Posidinia oceanica* (L.) Delile. *Bot. Mar.*, 32: 85–88.
- Bhadury, P. and P. C. Wright. 2004. Exploitation of marine algae: biogenic compounds for potential antifouling applications. *Planta*, 219: 561–578.
- Burgess, J. G., E. M. Jordan, M. Bregu, A. Mearns-Spragg and K. G. Boyd. 1999. Microbial antagonism: a neglected avenue of natural products research. *J. Biotech.*, 70:27–32.
- Caccamese, S., G. Azzolina, M. Furnari, M. Cormaci, and S. Grasso. 1980. Antimicrobial activity and antiviral activities of extracts from Mediterranean algae. *Bot. Mar.*, 23: 285–288.
- Caccamese, S., R. M. Toscano, G. Furnari and M. Cormaci. 1985. Antimicrobial activities of red and brown algae from southern Italy coast. *Bot. Mar.*, 28: 505–507.
- Chambers, L. D., K. R. Stokes, F. C. Walsh and R. J. K. Wood. 2006. Modern approaches to marine antifouling coatings. *Surf. Coat. Technol.*, 201: 3642–3652.
- Charaklis, W. G. 1981. Fouling biofilm development: A process analysis. *Biotechnol. Bioeng.*, 23: 1923–1960.
- Dalton, H. M., L. K. Poulsen, P. Halasz, M. L. Angles, A. E. Goodman and K. C. Marshall. 1994. Substratum-

- induced morphological changes in a marine bacterium and their relevance to biofilm structure. *J. Bacteriol.*, 176(22): 6900–6906.
- Davis, A. R., N. M. Targett, O. J. Mc Connell and C. Young. 1989. Epibiosis of marine algae and benthic invertebrates: natural products chemistry and other mechanisms inhibiting settlement and overgrowth. *Bioorg. Mar. Chem.*, 3: 86–114.
- De Nys, R., P. Steinberg, P. Willemsen, S. Dworjanyn, C. Gabelish and R. King. 1995. Broad spectrum effects of secondary metabolites from the red algae *Delisea pulchra* in antifouling assays. *Biofouling*, 8: 259–271.
- Fitton, J. H. 2006. Antiviral properties of marine algae. In: A. T. Critchley, M. Ohno, D. B. Largo (Eds.) *World Seaweed Resources*. Windows & Macintosh, ETI Information Services, Workingham, U.K., p. 7.
- Fletcher, M. 1980. Adherence of marine microorganisms to smooth surfaces. In: E. H. Beachey (Ed.) *Bacterial Adherence (Receptors and Recognition, series B)*. Chapman & Hall. London, 6: p. 347–371.
- Fletcher, R. 1989. A bioassay technique using the marine fouling green alga *Enteromorpha*. *Int Biodeterior. Biodegrad.*, 25: 407–422.
- Freile-Pelegrín, Y. and J. L. Morales. 2004. Antibacterial activity in marine algae from the coast of Yucatan, Mexico. *Bot. Mar.*, 47: 140–146.
- Ghosh, P., U. Adhikari, P. K. Ghosal, C. A. Pujol, M. J. Carlucci, E. B. Damonte and B. Ray. 2004. *In vitro* anti-herpetic activity of sulfated polysaccharide fractions from *Caulerpa racemosa*. *Phytochemistry*, 65: 3151–3157.
- Grossart, H. P., T. Kiørboe, K. Tang and H. Ploug. 2003. Bacterial colonization of particles: growth and interactions. *Appl. Environ. Microbiol.*, 69: 3500–3509.
- Hattori, T. and Y. Shizuri. 1996. A screening method for antifouling substances using spores of the fouling macroalga *Ulva conglobata* Kjellman. *Fish. Sci.* (Tokyo), 62: 955–958.
- Karande, A. A. 1968. Studies on marine fouling and boring organisms in Bombay harbour, *Proceedings of 2nd International Congress on Marine Corrosion and Fouling*. Athens, Greece. p. 563–569.
- Karande, A. A. 1978. Marine fouling and timber deterioration in sub oceanic Island of Andaman. *Indian J. Mar. Sci.*, 7: 39–43.
- Karande, A. A. and R. B. Srivastava. 1984. Pollution of coastal sea water and sulfate reducing bacteria, *Proceedings of the International Congress on Marine Corrosion and Fouling*, Ethens, Barcelon. p. 347–354.
- Langlois, G. 1975. Effect of algal exudates on substratum selection by motile telotrochs of the marine peritrich ciliate *Vorticella marina*. *J. Protozool.*, 22: 115–123.
- Mahasneh, I., M. Jamal, M. Kashasneh and M. Ziodeh. 1995. Antibiotic activity of marine algae against multiantibiotic resistant bacteria. *Microbiology*, 83: 23–26.
- Maki, J. 2002. Biofouling in the marine environment. In: H. Bitton (Ed.) *Encyclopedia of Environmental Microbiology*. Wiley & Sons. New York, p. 610–619.
- McLachlan, J. and J. S. Craigie. 1966. Antialgal activity of some simple phenols. *J. Phycol.*, 2: 133–135.
- Miki, W., K. Kon-ya and S. Mizobuchi. 1996. Biofouling and marine biotechnology: new antifoulants from marine invertebrates. *J. Mar. Biotechnol.*, 4: 117–120.
- Navqi, S., W. Solimabi, S. Kamat, L. Fernandes, C. Reddy, D. Bahkuni and B. Dhawan. 1980. Screening of some marine plants from the Indian coast for biological activity. *Bot. Mar.*, 24: 52–55.
- Padma Sridher, V., V. Lakshmi, H. Polassa, V. Santosh Reddy, C. H. Prasad Rao and G. Srimannarayana. 1984. Biological activity of some marine algal extracts. *Indian J. Mar. Sci.*, 1: 90–91.
- Padmini, S. R. P. 1991. Biological investigation of Indian marine algae and screening of some green, red and brown seaweeds for their antimicrobial activity. *Seaweed Research and Utilization*, 14(1): 37–43.
- Pesando, D. and B. Caram. 1984. Screening of marine algae from French Mediterranean coast for antibacterial and antifungal activity. *Bot. Mar.*, 27: 381–386.
- Philipps, D.W. and G. H. Towers. 1982. Chemical ecology of red algal bromophenols. I Temporal. Interpopulational and within thallus measurements of lanosol level in *Rhodomela larix* (Turnet) C. Agardh. *J. Exp. Mar. Bio. Ecol.*, 58: 285–293.
- Salvador, N., A. Gómez Garreta, L. Lavelli and M. A. Ribera. 2007. Antimicrobial activity of Iberian macroalgae. *Sci. Mar.*, 71: 101–113.
- Sastry, V.M.V.S. and G.R.K. Rao. 1994. Antibacterial substances from marine algae: successive extraction using benzene, chloroform and methanol. *Bot. Mar.*, 37: 357–360.

- Sieburth, J. M. and J. T. Conover. 1965. Sargassam tannin, an antibiotic which retards fouling. *Nature*, 208: 52–53.
- Srivastava, R. B., S. N. Gaonkar and A. A. Karande. 1990. Biofilm characterization in coastal waters of Bombay. *Proc., Indian Acad. Sci.*, 99: 163–173.
- Taskin, E., M. Ozturk and O. Kurt. 2001. Turkiye'nin yararlanilabilir denizel alg potansiyeli. XI. *Ulusal Su Urunleri Sempozyumu (4–6 Eylul 2001, Antakya)*, p. 225–232.
- Vlachos, V., A. T. Critchley and A. Von Holy. 1996. Establishment of a protocol for testing antimicrobial activity in southern African macroalgae. *Microbios.*, 88: 115–123.
- Wright, A. E. 1998. Isolation of marine natural products. In: J. P. Richard Cannell (Ed.) *Methods in Biotechnology*, Vol. 4 - *Marine Products Isolation*. Humana Press Inc., Totowa, NJ, USA, p. 365–408.

Received : 13/07/10

Accepted : 14/12/10